ONION THRIPS MANAGEMENT UPDATE

Zsofia Szendrei - vegetable entomologist **Adam Byrne** – research technician

Department of Entomology



New product registrations



syngenta[®]

Product now registered on onions in the U.S.:

Agri-Mek SC (abamectin)

Products in the pipeline:

- •Movento (spirotetramat) has federal label on many vegetable crops, but still not on onion (May 2013)
- •Benevia (cyantraniliprole) not labeled yet (target 2nd quarter 2013) @PND
- •Tolfenpyrad 15EC (tolfenpyrad) labeled on greenhouse crops, but not on onion (target 2014)







Section 18 for 2013:

Movento





Goals in 2012:

- compare thresholds to weekly sprays
- determine Radiant rate
- test experimental products in programs

Methods:

- •All treatments included non-ionic surfactant (Dyne-Amic, 0.5% v/v)
- •50 gallons per acre, 40 psi, single nozzle boom
- •counted thrips weekly on 10 randomly selected plants per plot
- visual rating of thrips damage on two dates
- •first insecticide application was on Jun 20, 2012
- •three-row beds, 15 ft long plots

Which products were included in the 2012 MSU onion thrips insecticide study?

Chemical Class	Products used in onion thrips insecticide trial	
pyrethroid	-	Group 3A
organophosphate	-	Group 1B
neonicotinoid	-	Group 4A
carbamate	Lannate	Group 1A
spinosyn	Radiant	Group 5
abamectin	Agri-Mek	Group 6
pyrazol	Tolfenpyrad ¹	Group 21A
spirotetramat	Movento ²	Group 23
cyantraniliprole	Benevia ¹	Group 28

¹ experimental product

² can only be used under Section 18 Exemption

















dark green bars – sequence started with Movento

light green bars – sequence started with Movento, used thresholds blue bars – sequence didn't start with Movento or had no Movento




Radiant - max. 30 fl oz per season







Can *Torac* replace *Movento*?



Does the position of *Benevia* matter in the program?




Can using thresholds work as well as weekly sprays?




1: Treatments 10 vs. 11

threshold (Radiant – 3 thrips/leaf, all other products – 1 thrips/leaf) no threshold, weekly applications




2: Treatments 12 vs. 13

threshold (Radiant – 3 thrips/leaf, all other products – 1 thrips/leaf) no threshold, weekly applications




3: Treatments 14 vs. 15


threshold (Radiant – 3 thrips/leaf, all other products – 1 thrips/leaf) no threshold, weekly applications



Blue lines indicate treatments where Movento was applied at the beginning of the season



Using thresholds and new products



Yields

- •average yield across treatments: 726.8 cwt/acre
- •all insecticide programs yielded significantly more than the untreated control
- •all insecticide programs (except 14) had more jumbos than untreated control
- •jumbos weighed significantly more in all insecticide programs (except 14, 5) compared to control



Research Plans for 2013

- 1. Demonstrate effective insecticide rotations that take into account the principles of insecticide resistance management.
- 2. Test experimental insecticide in program.
- 3. Continue to develop thresholds for registered and new insecticides.

Acknowledgements

- Krummrey Farms
- •The Michigan Onion Growers Association
- •The undergraduate students who spent hours counting thrips in the field
- •Bayer CropScience, DuPont, Nichino America, Dow AgroSciences, Syngenta Crop Protection